

# Cefditoren

## Antibiotic Class:

Third-Generation Cephalosporin

## Antimicrobial Spectrum:

*Staphylococcus aureus* (methicillin susceptible), Coagulase negative Staphylococci, *Streptococcus pneumoniae* (penicillin susceptible), *Streptococcus spp.*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Enterobacteriaceae*, *E. coli*

## Mechanism of Action:

Cephalosporins exert bactericidal activity by interfering with bacterial cell wall synthesis and inhibiting cross-linking of the peptidoglycan. The cephalosporins are also thought to play a role in the activation of bacterial cell autolysins which may contribute to bacterial cell lysis.

## Pharmacodynamics:

Cephalosporins exhibit time-dependent killing ( $T > MIC$ )

## Pharmacokinetics:

Dose of 200mg: Cmax: 3.4 mcg/L; Tmax: 2.0 hour; Half-life: 1.1 hours; Table 10

## Adverse Effects:

Hypersensitivity: Maculopapular rash, Urticaria, Pruritis, Anaphylaxis/angioedema, eosinophilia

Hematologic: Hypoprothrombinemia, Neutropenia, Leukopenia, Thrombocytopenia

GI: Diarrhea, *C. difficile* disease

Renal: Interstitial nephritis

Table 14

## Dosage:

PO: 200mg tablets

## Dosing in adults:

Acute exacerbation of chronic bronchitis: 400 mg PO q12h x 10 days

Pharyngitis: 200mg PO q12h x 10 days

Community acquired pneumonia: 400 mg PO q12h x 14 days

Tonsillitis: 200mg PO q12h x 10 days

## Dosing in pediatrics:

Not recommended

## Disease state based dosing:

Renal failure: CrCl 30-49 mL/min: 200 mg q12h

CrCl < 30 mL/min: 200 mg q24h

Hepatic failure: No dosing changes recommended at this time.

**Contraindications/Warnings/Precautions:**

Contraindications: Carnitine deficiency or inborn errors of metabolism that may result in significant carnitine deficiency, Milk protein hypersensitivity

Precautions: hypersensitivity to penicillins, history of gastrointestinal disease, particularly colitis, renal impairment

**Drug Interactions:**

Antacids: decreased cefditoren effectiveness

Cimetidine: decrease cefditoren serum concentrations

Famotidine: decrease cefditoren serum concentrations

Live Typhoid Vaccine: decreased immunological response to the typhoid vaccine

Nizatidine: decrease cefditoren serum concentrations

Probenecid: increased cefditoren serum concentrations and bioavailability

Ranitidine: decrease cefditoren serum concentrations

**Pregnancy:**

Category B: No evidence of risk in humans but studies inadequate.

**Monitoring Requirements:**

Therapeutic: Culture and sensitivities, serum levels, signs and symptoms of infection, white blood cell count

Toxic: Urinalysis, BUN, SCr, AST and ALT, skin rash, Neutropenia and leukopenia,

Prothrombin time in patients with renal or hepatic impairment or poor nutritional state, as well as patients receiving a protracted course of antimicrobial therapy, and patients previously stabilized on anticoagulant therapy.

**Brand names/Manufacturer:** Spectracef®/Purdue